

564/387

Europäisches Patentamt  
European Patent Office  
Office européen des brevets

⑪ Publication number:

0 000 896  
A2

⑫

## EUROPEAN PATENT APPLICATION

⑰ Application number: 78100611.9

⑱ Date of filing: 07.08.78



⑲ Int. Cl.2: C 07 C 87/28, C 07 C 85/02,  
C 07 C 85/24, C 07 C 87/29,  
C 07 C 87/36, C 07 C 87/40,  
C 07 C 87/45, C 07 C 101/28,  
C 07 D 203/08, C 07 D 207/06,  
C 07 D 207/32  
// (C07D401/06, 211/14,  
213/38), (C07D409/04, 211/20,  
333/58)

⑳ Priority: 19.08.77 CH 10203/77  
19.08.77 CH 10202/77  
24.10.77 CH 12910/77  
24.10.77 CH 12909/77

㉑ Applicant: SANDOZ LTD., Lichtstrasse 35, CH-4002  
Basel (CH)

㉒ Date of publication of application: 07.03.79  
Bulletin 79/5

㉓ Inventor: Hamberger, Helmut, Dr., 4-5 Josef-  
Kyrlegasse, A-1130 Vienna (AT)  
Inventor: Stephen, Adrian, Dr., 8 Jesuitensteig, A-1238  
Vienna (AT)  
Inventor: Stütz, Anton, Dr., 113/7 Donaustasse,  
A-2344 Maria Enzersdorf (AT)  
Inventor: Stütz, Peter, Dr., 15 Nöstbergergasse,  
A-1130 Vienna (AT)

㉔ Designated Contracting States: BE CH DE FR GB LU  
NL SE

㉕ Propenyl amines, processes for their production and pharmaceutical compositions containing them.

㉖ The present invention provides propenylamines useful  
as anti-mycotic agents.


934.772

EP 0 000 896 A2

PROPENYL - AMINES, PROCESSES FOR THEIR PRODUCTION AND  
PHARMACEUTICAL COMPOSITIONS CONTAINING THEM

This invention relates to propenyl-amines, processes for their production and pharmaceutical compositions containing them.

The present invention provides a compound of  
5 formula I,



wherein a) (1)  $R_1$  is a radical of formula IIa,



wherein  $R_7$  and  $R_8$ , independently, are hydrogen, halogen of atomic number from 9 to 53, trifluoromethyl, hydroxy, nitro, lower alkyl or lower alkoxy, or a radical of formula IIb, IIc, IId, IIe,


0000896

- 1 -

900-9192

What are claims:


1. A compound of formula I,




wherein a) (i)  $R_1$  is a radical of formula IIa,




wherein  $R_7$  and  $R_8$ , independently, are hydrogen, halogen of atomic number from 9 to 53, trifluoromethyl, hydroxy, nitro, lower alkyl or lower alkoxy, or a radical of formula IIb, IIc, IIId, IIe,




IIb



IIc



IIId



IIe

wherein  $R_9$  is hydrogen, halogen of atomic number from 9 to 53, hydroxy, lower alkyl or lower alkoxy,

X is oxygen, sulphur, imino, lower alkylimino or a radical of formula


$-(CH_2)_r-$  wherein r is 1, 2 or 3,

s is 3, 4 or 5, and


t is 2, 3 or 4, and

$R_2$  is hydrogen or lower alkyl, or

(ii)  $R_1$  and  $R_2$  together with the carbon atom to which they are bound form a radical of formula IIIf or IIg,



IIIf



IIg

wherein p is 1, 2 or 3,

$R_3$  and  $R_5$ , independently, are hydrogen or lower alkyl,

$R_4$  is alkyl ( $C_{1-6}$ ), alkenyl ( $C_{3-12}$ ),  
alkynyl ( $C_{3-12}$ ) or cycloalkyl ( $C_{3-8}$ )-  
alkyl ( $C_{1-6}$ ); and

$R_6$  is (i) an aromatic, five-membered  
heterocycle containing one oxygen,  
sulphur or nitrogen hetero-ring atom  
and optionally an additional one or two  
nitrogen hetero-ring atoms and being  
optionally substituted on a carbon ring  
atom by halogen of atomic number from  
9 to 53, hydroxy, lower alkyl or lower  
alkoxy, and any nitrogen ring atom  
present being optionally substituted  
when possible, by lower alkyl, (ii) a  
radical of formula IIIa,




wherein  $R_9$  is as defined above,  
(iii) a radical of formula IIIb,



wherein  $R_{10}$  is alkyl ( $C_{1-12}$ ), alkenyl  
( $C_{3-12}$ ), alkynyl ( $C_{3-12}$ ) cycloalkyl ( $C_{3-8}$ )  
alkyl ( $C_{1-6}$ ), phenyl-alkyl ( $C_{7-12}$ ),  
phenyl, phenylalkoxy ( $C_{7-16}$ ), or amino-  
alkyl ( $C_{1-12}$ );

(iv) a radical of formula IIIc, IIIId or IIIe,




wherein  $R_{11}$ ,  $R_{12}$  and  $R_{13}$ , independently, are hydrogen or lower alkyl,

$m$  is a whole number from 0 to 4,

$n$  is a whole number from 0 to 3, and

$v$  is a whole number from 0 to 5,

(v) a radical of formula IIIf,




wherein  $R_{14}$  is lower alkyl, alkoxy ( $C_{1-12}$ )-carbonyl, alkenyl ( $C_{3-12}$ ), alkynyl ( $C_{3-12}$ ), phenylalkyl ( $C_{7-12}$ ) or phenyl,

$R_{15}$  and  $R_{16}$ , independently, are hydrogen or lower alkyl, and

$q$  is a whole number from 0 to 5, or

(vi) a radical of formula IIIg



IIIg

5

wherein R<sub>17</sub> and R<sub>18</sub>, independently, are hydrogen, halogen of atomic number from 9 to 53, trifluoromethyl, hydroxy, nitro, lower alkyl or lower alkoxy, with the proviso that one of R<sub>17</sub> and R<sub>18</sub> is other than hydrogen, and with the general proviso that R<sub>1</sub> is not a radical of formula IIa when R<sub>6</sub> is a radical of formula IIIg or phenyl, R<sub>2</sub> is hydrogen and R<sub>3</sub> is hydrogen or lower alkyl,

10

b) R<sub>1</sub> is a radical of formula IIa to IIe, as defined above,

R<sub>2</sub>, R<sub>5</sub> and R<sub>6</sub> are as defined above, and

R<sub>3</sub> and R<sub>4</sub> together are -(CH<sub>2</sub>)<sub>u</sub> - wherein u

15

is a whole number from 1 to 8.

Any lower alkyl or lower alkoxy radical has preferably 1 to 4 carbon atoms, especially 2 or 1 carbon atoms. Any alkyl(C<sub>1-12</sub>) moiety is preferably alkyl(C<sub>2-8</sub>); phenyl-alkyl or phenylalkoxy has preferably 7 carbon atoms.

20

Any alkenyl or alkynyl radical has preferably 3 to 6 carbon atoms, especially 3 or 4 carbon atoms. Preferably the multiple bond is in other than the α, β position and is conveniently in the remote terminal position. An example of an alkenyl group is allyl. An example of an alkynyl group is

propinyl. Cycloalkylalkyl has preferably an alkyl moiety of 1 to 4 carbon atoms, especially 2 or 1 carbon atoms, and a cycloalkyl moiety preferably of 3 to 6 carbon atoms.

When  $R_4$  is cycloalkylalkyl this is especially cyclopentyl alkyl or cyclohexylalkyl. When  $R_{10}$  is cycloalkylalkyl this is especially cyclopropylalkyl or cyclobutylalkyl.

Conveniently  $R_7$  and  $R_8$  are identical and are both hydrogen. Conveniently  $R_9$  is hydrogen or halogen. In IIb and IIc the bond to the carbon atom to which  $R_2$  and  $R_3$  are attached is conveniently attached meta to X and para to the ring nitrogen, respectively. X is conveniently sulphur, imino or lower alkylamino.  $R_1$  is preferably a radical of formula IIb, IIc or IId or especially IIa. —————  
—————  $R_2$  is preferably hydrogen.  $R_3$  is preferably hydrogen and  $R_4$  is conveniently alkyl.  $R_5$  is conveniently hydrogen.  $R_6$ , when it is a heterocycle, conveniently contains one oxygen or sulphur atom or one or two nitrogen atoms. Preferably the bond linking  $R_6$  to the vinylene moiety is attached to a ring carbon atom adjacent to a ring heteroatom. Conveniently the ring is unsubstituted or substituted by lower alkyl.  $R_{10}$  is conveniently phenylalkoxy. IIa is conveniently optionally substituted 2 or 4-pyridyl. In IIIc, IIId, IIIe it is to be appreciated that the bond linking  $R_6$  to the vinylene moiety and  $R_{11}$  to  $R_{13}$  may be attached to any of the ring carbon atoms present.

IIIc is preferably a cycloalk-1-en-1-yl radical. Preferably  $R_{11}$  to  $R_{13}$  are hydrogen. q is conveniently 0 or 1.



Any double bond in IIIf is conveniently trans.  $R_{14}$  is conveniently alkoxy ( $C_{1-8}$ ) carbonyl, phenyl or alkyl or phenalkyl.  $R_{17}$  is conveniently halogen and  $R_{18}$  is conveniently hydrogen.  $R_6$  is conveniently IIIc. u is conveniently 3, 4 or 5, more conveniently 4.

The values m, n, p, q, s, t and v are conveniently chosen to produce a five or six-membered ring.

The double bond between  $R_6$  and the nitrogen atom preferably has the trans configuration.


10 Halogen is conveniently fluorine, or preferably bromine or chlorine.

When  $R_1$  is IIb or IIe and  $R_6$  is IIIa it is to be appreciated that the two radicals  $R_9$  may be the same or different.

15 The present invention also provides a process for the production of a compound of formula I, which comprises  
a) reacting a compound of formula IV,




wherein  $R_1$  to  $R_4$  are as defined above,  
with a compound of formula V,



20 wherein A is a leaving group, and

$R_5$  and  $R_6$  are as defined above, or

b) producing a compound of formula Ia,




wherein  $R_1$  to  $R_3$  and  $R_5$  are as defined above, and

$R_4^I$  and  $R_6^I$  are as defined above for  $R_4$  and  $R_6$

5 respectively, with the proviso that they each are other than alkynyl,


by reducing a compound of formula VI,



wherein  $R_1$  to  $R_3$ ,  $R_4^I$ ,  $R_5$  and  $R_6^I$  are as defined above,


or

10 c) producing a compound of formula Ib,




wherein  $R_1$  to  $R_6$  are as defined above,

by isomerising photochemically a compound of formula Ic,



wherein  $R_1$  to  $R_6$  are as defined above, or

d) producing a compound of formula Id,




wherein  $R_1$ ,  $R_2$ ,  $R_5$  and  $R_6$  are as defined above,

$R_3^I$  is hydrogen or lower alkyl,

5  $R_4^{II}$  is alkyl( $C_{1-6}$ ), alkenyl( $C_{3-12}$ ), alkynyl  
( $C_{3-12}$ ) or cycloalkyl( $C_{3-8}$ )alkyl( $C_{1-6}$ );

by introducing the group  $R_4^{II}$  into a compound of formula VII,



wherein  $R_1$ ,  $R_2$ ,  $R_3^I$ ,  $R_5$  and  $R_6$  are as defined above.

Process a) may be effected in conventional manner  
10 for the production of tertiary amines by condensation from  
analogous starting materials. The process may be effected  
in an inert solvent such as a lower alkanol, e.g. ethanol,

optionally in aqueous admixture, an aromatic hydrocarbon solvent, e.g. benzene or toluene, a cyclic ether, e.g. dioxane or a carboxylic acid dialkylamide solvent, e.g. dimethylformamide. The reaction temperature is conveniently from room temperature to the boiling temperature of the reaction mixture, preferably room temperature. The reaction is conveniently effected in the presence of an acid binding agent, such as an alkali metal carbonate, e.g. sodium carbonate. The leaving group A is conveniently iodine or preferably chlorine or bromine, or an organic sulphonyloxy group having 1 to 10 carbon atoms, e.g. alkylsulphonyloxy, preferably having 1 to 4 carbon atoms such as mesyloxy, or alkylphenylsulphonyloxy preferably having 7 to 10 carbon atoms such as tosyloxy.

Process b) may be effected in conventional manner for catalytic hydrogenation in order to produce a compound of formula Ia wherein the double bond adjacent to  $R_6^I$  has the cis configuration. Alternatively, the process may be effected in conventional manner for a complex metal hydride reduction in order to produce a compound of formula Ia wherein the double bond has the trans configuration.

The catalytic hydrogenation may be effected in a solvent, e.g. methanol, ethanol, methylene chloride, pyridine or ethyl acetate. The catalyst is preferably palladium on a carrier material such as  $BaSO_4$  or  $CaCO_3$ . The catalyst

may be pretreated, e.g. with a lead salt, so as to be partially poisoned (e.g. a Lindlar catalyst). The hydrogenation may be effected at room temperature and at normal pressure.

5           The metal hydride reduction may be effected in conventional manner for a lithium aluminium hydride or a diisobutylaluminium hydride reduction. The reduction is conveniently effected in an inert solvent such as toluene or benzene. The reaction is conveniently effected at room  
10 temperature.

Process c) may be effected in conventional manner for a photochemical isomerisation of a cis alkene. The reaction may be effected in a solvent such as benzene, petroleum ether, ethanol, or preferably cyclohexane. The  
15 solution is conveniently irradiated with light from a mercury high or low pressure lamp. The reaction is conveniently effected at room temperature. If desired, an appropriate sensitizer such as eosine or a catalyst such as diphenyldisulphide may be present.

20           Process d) may be effected in manner conventional for the "alkylation" of secondary amines (the term "alkylation" being used here to denote introduction of any of the hydrocarbyl groups  $R_4^{II}$ ), for example by direct "alkylation" with an "alkylating" agent, for example a halide or sulphate,


or by reductive alkylation, in particular by reaction with an appropriate aldehyde and subsequent or simultaneous reduction. Reductive "alkylation" is suitably effected in an inert organic solvent, such as a lower alkanol, e.g. 5 methanol, and at an elevated temperature, in particular at the boiling temperature of the reaction medium. The subsequent reduction may be effected with, for example, a complex metal hydride reducing agent, e.g.  $\text{NaBH}_4$  or  $\text{LiAlH}_4$ . The reduction may also be effected simultaneously to the 10 alkylation, for example by use of formic acid which may serve both as reducing agent and as a reaction medium.

It is to be appreciated that in any of the above processes, side reactions may occur, e.g. reduction of halogen to hydrogen, reduction of a nitro group to an amino 15 group, reduction of an alkenyl moiety to an alkyl moiety and/or reduction of a keto moiety to a carbinol moiety in processes b) or process d) when reductive alkylation is used, or simultaneous cis/trans isomerisation of any double bond present in  $\text{R}_4$  or  $\text{R}_6$  when process c) is used. The reaction 20 conditions should be chosen to avoid such side reactions, and the desired final product isolated using conventional purification techniques, e.g. thin layer chromatography.

Free base forms of the compounds of formula I may be converted into salt forms and vice versa. Suitable acids


for acid addition salt formation include hydrochloric acid, fumaric acid and naphthalene-1,5-disulphonic acid.

The starting materials are either known or may be made in conventional manner. For example non-cyclic amines of formula IV may be made by condensing a compound of formula VIII,



or the corresponding iodide or chloride, with a compound of formula  $\text{R}_4^{\text{II}}\text{NH}_2$ .


The cyclic amines of formula IV may be made as follows:-



wherein Alk = lower alkyl.

The compounds of formula VI are new and may be made by reacting an appropriate amine of formula IV with compounds of formulae  $\text{R}_5\text{-CHO}$  and  $\text{HC} \equiv \text{CR}_6^{\text{I}}$  under Mannich reaction conditions.

The compounds of formula VII are also new and may be made as follows:-



In the following Examples all temperatures are uncorrected and in degrees Centigrade.

5 In the tables hereinafter, the following indications are used:-

- 1) All double bonds have the trans configuration; all alkyl groups are unbranched unless stated otherwise.
- 2) If no melting point is given, the free base form of the  
10 compound is obtained and this is an oil. Melting points are for the free base form unless specified otherwise.
- 3) Monohydrochloride salt form.
- 4) Dihydrochloride salt form.



EXAMPLE 1: 4-[N-methyl-N-(1-naphthylmethyl)]aminocrotonic  
acid ethyl ester [process a)]

1.9 g of bromocrotonic acid ethyl ester are added dropwise to a mixture of 1.7 g of N-methyl-N-(1-naphthyl-  
5 methyl)amine, 1.4 g of  $K_2CO_3$  and 10 ml dimethylformamide. After the mixture is stirred for 18 hours at room temperature, it is filtered and evaporated under a vacuum. The residue is chromatographed on silica-gel using benzene/ethyl acetate (1:1) as solvent to yield the title compound in  
10 free base form, as an oil, after evaporating the appropriate fractions.

The title compound may also be made in analogous manner to Examples 3, 4 and 5.

EXAMPLE 2: N-(3-cyclohex-1-en-1-yl-2-cis-propenyl)-N-  
15 methyl-N-(1-naphthylmethyl)amine [process b)]

5 g of N-(3-cyclohex-1-en-1-yl-propynyl)-N-methyl-N-(1-naphthylmethyl)amine are hydrogenated in absolute pyridine using 750 mg  $Pd/BaSO_4$  as catalyst at room temperature and normal pressure, until the calculated amount of  
20 hydrogen is taken up. The reaction mixture is filtered and the pyridine removed in a vacuum. The residue is chromatographed on silica-gel using benzene/ethylacetate (9:1) to yield the title compound in free base form as an oil after evaporating the appropriate fractions,

m.p. (hydrochloride) 184-188°.

The title compound may also be made in analogous manner to Examples 1 and 5.

5 EXAMPLE 3: N-(3-cyclohexyl-2-trans-propenyl)-N-methyl-N-(1-naphthylmethyl)amine [process b]

28 ml of a 1.2 molar solution of diisobutylaluminum hydride in toluene are added to 5 g of N-(3-cyclohexyl-propynyl)-N-methyl-N-(1-naphthylmethyl)amine in absolute benzene. After the mixture is stirred for 3 hours at 40°, 10 water is carefully added. The organic phase is separated off, dried and evaporated to yield the title compound in free base form, as an oil.

The title compound may also be prepared by following Examples 1, 4 and 5.

15 EXAMPLE 4: N-(3-cyclohex-1-en-1-yl-2-trans-propenyl)-N-methyl-N-(1-naphthylmethyl)amine [process c]

1.2 g of N-(3-cyclohex-1-en-1-yl-2-cis-propenyl)-N-methyl-N-(1-naphthylmethyl)amine are irradiated for 3 hours with a Hg high pressure lamp ( $\lambda > 300$  nm) in 1 litre 20 cyclohexane in the presence of 50 mg diphenyldisulphide at room temperature under an inert gas atmosphere. After the solvent is evaporated, the title compound is obtained in free base form and converted into the hydrochloride, m.p.

184-188°

The title compound may also be prepared by following Examples 1, 3 and 5.

EXAMPLE 5: N-methyl-N-[3-(5'-methyl-2'-thienyl)-2-trans-

5 propenyl)-N-(1-naphthylmethyl)amine [process d)]


a) 15.2 g of 3-(5'-methyl-2'-thienyl)prop-2-enal and 15.7 g of 1-aminomethylnaphthalene in 350 ml benzene are boiled under reflux until the calculated amount of water has boiled off. 3.6 g of the resulting Schiff base in 100 ml methanol  
10 are boiled under reflux with 5 g  $\text{NaBH}_4$  for 30 minutes to yield N-[3-(5'-methyl-2'-thienyl)-2-trans-propenyl)]-N-(1-naphthylmethyl)amine, which is used directly in the next stage. [To isolate this intermediate the reaction mixture is evaporated in a vacuum; the residue is partitioned between  
15 aqueous sodium carbonate solution and diethyl ether and the organic phase is evaporated}.

b) The crude reaction mixture obtained in step a) is treated with 20 ml 37% aqueous formaldehyde solution. The mixture is boiled under reflux for 60 minutes, subjected to ice-cooling,  
20 treated with 9 g  $\text{NaBH}_4$  and stirred for another 60 minutes at room temperature. The mixture is evaporated in a vacuum to a residue which is partitioned between aqueous  $\text{NaHCO}_3$  and diethyl ether. The organic phase is dried and evaporated

to yield the title compound in free base form as an oil,  
m.p. (hydrochloride) 140-156°.


The title compound may also be prepared in analogous  
manner to Examples 1, 3 and 4.





5 In analogous manner to that described in Examples  
1, 3, 4 and 5, the following trans compounds of formula Ie  
may be produced:




Ie

wherein  $R_1$  and  $R_6$  are as follows:

10	Ex.	$R_1$	$R_6$ <sup>1)</sup>	m.p. <sup>2)</sup>
	6	1-naphthyl	-COOC <sub>5</sub> H <sub>11</sub>	
	7	1-naphthyl	-COOC <sub>8</sub> H <sub>17</sub>	
	8	1-naphthyl	-COOCH <sub>2</sub> C <sub>6</sub> H <sub>5</sub>	
	9	1-naphthyl	2-thienyl	182-187 <sup>3)</sup>
15	10	1-naphthyl	2-furyl	
	11	1-naphthyl		175-185 <sup>4)</sup>
	12	1-naphthyl	Cyclohex-3-en-1-yl	
	13	1-naphthyl	2-pyridyl	
	14	1-naphthyl	3-pyridyl	
20	15	1-naphthyl	4-pyridyl	174-178 <sup>4)</sup>
	16	1-naphthyl	-CH=CH.C <sub>6</sub> H <sub>5</sub>	170-174 <sup>3)</sup>

Ex.	R <sub>1</sub>	R <sub>6</sub> <sup>1)</sup>	m.p. <sup>2)</sup>
17	1-naphthyl	-CH=CH.C <sub>4</sub> H <sub>9</sub>	
18	1-naphthyl	2-pyrrolyl	
19	1-naphthyl	Cyclohept-1-en-1-yl	193-196 <sup>3)</sup>
20	1-naphthyl	Cyclopent-1-en-1-yl	
21	1-naphthyl		180-184 <sup>3)</sup>
22	1-naphthyl	-CH=CH-COOC <sub>2</sub> H <sub>5</sub>	
23	1-tetralinyl	C <sub>6</sub> H <sub>5</sub>	
24	1-tetralinyl	Cyclohex-1-en-1-yl	
25	5-tetralinyl	C <sub>6</sub> H <sub>5</sub>	
26	4-quinolyl	C <sub>6</sub> H <sub>5</sub>	
27	3-benzo[b]thienyl	C <sub>6</sub> H <sub>5</sub>	
28	3-benzo[b]thienyl	Cyclohex-1-en-1-yl	175-177 <sup>3)</sup>
29	1-naphthyl		
30	1-naphthyl		
31		C <sub>6</sub> H <sub>5</sub>	

In analogous manner to that described above for Examples 1, 3 and 4 there may be produced the following trans compounds of formula If,



wherein  $R_1$ ,  $R_6$  and  $u$  are as follows:

Ex.	$R_1$	$R_6^{1)}$	$u$	m.p. $^{2)}$
32	1-naphthyl	$C_6H_5$	4	203-205 <sup>3)</sup>
33	1-naphthyl	$C_6H_5$	3	75-78
34	1-naphthyl	$C_6H_5$	5	171-175 <sup>3)</sup>
35	1-naphthyl	$COOC_5H_{11}$	4	
36	1-naphthyl	$C_6H_5$	1	
37	1-naphthyl	Cyclohex-1-en-1-yl	4	
38	1-naphthyl	4-F- $C_6H_4$	4	
39	1-naphthyl	3-pyridyl	4	
40	3-benzo[b]thienyl	$C_6H_5$	4	
41	5-tetralinyl	$C_6H_5$	4	150-155 <sup>3)</sup>

EXAMPLE 42:

In analogous manner to that described in Examples 1 and 2, the following cis compound of formula I may be produced:

- 5 aa) N-(3-cyclohex-1-en-1-yl-2-cis-propenyl)-2-(1'-naphthyl)-piperidine ; free base-oil.

EXAMPLES 43-47:

In analogous manner to that described in Examples 1, 3, 4 and 5, the following compounds of formula I may be  
10 produced:

- 43) N-cinnamyl-N-methyl-N-[2-(1'-naphthyl)-2-propyl]amine;  
free base-oil;
- 44) N-(1-acenaphthenyl)-N-methyl-N-(3-phenyl-2-trans-propenyl)amine, m.p. (hydrochloride) 210-216°;
- 15 45) N-(1-acenaphthenyl)-N-methyl-N-[3-(5'-methyl-2'-thienyl)-2-trans-propenyl]amine, free base-oil;
- 46) N-(6,7,8,8a-tetrahydro-1-acenaphthenyl)-N-methyl-N-(3-phenyl-2-trans-propenyl)amine, m.p. (hydrochloride) 185-192°;
- 20 47) N-methyl-N-(2,3-dihydro-1-phenalenyl)-N-(3-phenyl-2-trans-propenyl), free base-oil.

NMR data on the above-mentioned compounds of formula I, obtained as oils, are given in the following table. The data comprises peak position in ppm relative to TMS as standard in  $\text{CDCl}_3$ ; type of peak (D = doublet; DD = double  
25

doublet; DT = double triplet; M = multiplet; Q = quartet; S = singlet; T = triplet) and in parentheses the corresponding number of hydrogen atoms.

Ex.	NMR Data
1	1.2 T (3); 2.25 S (3); 3.2 M (2); 3.9 S (2); 4.2 Q (2); 6.0 D (1); 6.85 - 7.2 M (1); 7.3 - 7.9 M (6); 8.2 (1).
3	0.8 - 2.2 (11); 2.2 S (3); 3.0 M (2); 3.85 S (2); 5.6 M (2); 7.3 - 7.9 (6); 8.3 M (1)
6	0.7 - 1.8 M (9); 2.25 S (3); 3.2 M (2); 3.9 S (2) 4.1 T (2); 6.0 D (1); 6.9 - 7.2 M (1); 7.3 - 7.9 M (6); 8.3 M (1); 4.1 T (2).
7	0.8 - 1.8 M (15); 2.3 S (3); 3.25 M (2); 3.95 S (1); 4.1 T (2); 6.0 M (1); 6.9 - 7.2 M (1); 7.4 - 7.9 M (6); 8.2 - 8.4 M (1)
8	2.3 S (3); 3.2 M (2); 3.9 S (2); 5.15 S (2); 6.1 M (1); 6.9 - 7.9 M (13); 8.2 - 8.4 M (1).
10	2.28 S (3); 3.28 T (2); 3.94 S (2); 6.25 M (4); 7.42 M
12	1 - 2.5 M (7); 2.2 S (3); 3.0 M (2); 3.85 S (2); 6.4 - 6.8 M (4); 7.2 - 7.9 (6); 8.2 - 8.4 M (1).
13	2.3 S (3); 3.35 D (2); 4.0 S (2); 7.0 - 8.0 M (9); 8.2 - 8.4 M (1); 8.55 M (1)
14	2.3 S (3); 3.25 D (2); 3.95 S (2); 6.2 - 6.7 M (2); 7.0 - 7.8 M (8); 8.2 - 8.7 M (3)



Ex.	NMR Data
17	0.8 - 1.5 (7); 3.0 - 3.3 M (2); 3.2 S (3); 3.15 D (2); 3.9 S (2); 5.5 - 6.4 M (4); 7.2 - 8.4 (7).
18	2.26 S (3); 3.22 D (2); 3.94 S (2); 5.85 DT; 6.18 M (2); 6.45 D (1); 6.74 M (1); 7.4 - 8.5 M (6); 8.3 M (1)
20	1.7 - 2.1 (2); 2.2 - 2.6 (4); 2.2 S (3); 3.15 D (2); 3.86 S (2); 5.5 - 5.85 M (2); 6.45 D (1); 7.2 - 7.9 (6); 8.3 M (1).
22	1.28 T (3); 2.24 S (3); 3.2 D (2); 3.9 S (2); 4.2 Q (2); 5.84 D (1); 6.2 - 6.5 M (2); 7.2 - 7.9 (7); 8.2 - 8.3 M (1)
23	1.6 - 1.9 M (4); 2.2 S (3); 2.7 - 2.9 M (4); 3.2 D (2); 3.45 S (2); 6.1 - 6.7 M (2); 6.9 - 7.5 (8)
24	1.4 - 1.9 (8); 2.0 - 2.3 (4); 2.17 S (3); 2.8 M (4); 3.06 D (2); 3.4 S (2); 5.4 - 5.8 (2); 6.16 D (1); 6.9 - 7.2 (4)
25	1.6 - 2.1 M (4); 2.35 S (3); 2.2 - 3.4 M (7); 6.0 - 6.6 M (2); 7.0 - 7.4 M (9)
26	2.3 S (3); 3.25 D (2); 3.9 S (2); 6.2 - 6.7 M (2); 7.1 - 7.8 (9); 8.2 M (2); 8.85 D (1)
27	2.3 S (3); 3.25 D (2); 3.8 S (2); 6.2 - 6.7 M (2); 7.2 - 7.5 (8); 7.8 - 8.0 M (2)
29	1.1 D (3); 1.3 - 2.8 (7); 1.75 S (3); 2.25 S (3); 3.25 D (2); 3.91 M (2); 5.8 DT (1); 6.5 D (1); 7.3 - 7.9 (6); 8.3 M (1)

Ex.	NMR Data
30	1.33 T (3); 2.27 S (3); 3.26 D (2); 3.91 Q (2); 3.94 S (2); 6.0 - 6.7 M (5); 7.3 - 7.7 M (4); 7.7 - 7.9 M (2); 8.3 M (1);
31	2.28 S (3); 3.02 S (3); 3.18 D (2); 3.86 S (2); 6.2 - 6.6 M (2); 6.65 (1); 7.0 - 7.4 M (8); 8.0 M (1)
35	2.5 - 2.8 DD (1); 3.1 - 3.5 M (3); 3.8 - 4.0 M (1); 4.1 T (2); 5.9 D (1); 6.7 - 7.05 M (1); 7.3 - 8.0 M (6); 8.5 M (1)
36	1.9 - 2.1 M (2); 2.95 - 3.1 M (1); 3.4 D (1); 6.2 - 6.8 M (2); 7.0 - 8.4 (12);
37	2.4 - 2.7 DD (1); 3.0 - 3.5 M (3); 3.7 - 4.0 (1); 5.2 - 5.6 M (3); 5.9 D (J = 16Hz) (1); 7.3 - 8.0 (6); 8.3 - 8.8 (1)
38	1.3 - 2.4 M (7); 2.65 DD (1); 3.15 - 3.25 M (2); 3.65 - 3.95 (1); 5.8 - 6.4 (2); 6.8 - 7.9 (10); 8.3 - 8.8 broad S (1)
39	1.2 - 2.8 (8); 3.15 - 3.45 (2); 3.8 - 4.0 (1); 6.2 M (2); 7.0 - 7.9 (8); 8.2 - 8.6 (3);
40	1.2 - 2.4 (7); 2.6 - 2.85 DD; 3.15 - 3.8 (3); 6.1 - 6.8 (2); 7.1 - 7.6 (8); 7.8 - 7.9 M (1); 8.15 - 8.25 M (1)
42	2.8 - 3.1 DD (1); 3.1 - 3.5 M (3); 3.7 - 4.0 (1); 5.2 - 5.6 M (3); 5.8 D (J = 13Hz) (1); 7.3 - 8.0 (6); 8.3 - 8.8 (1)

0000896

- 25 -

900-9192

Ex.	NMR Data
43	1.62 S (6); 2.34 S (3); 3.15 D (2); 6.03 DT (1); 6.40 D (1); 7.1 - 7.9 (11); 9.55 M (1)
45	2.18 S (3); 2.40 S (3); 3.20 M (4); 4.95 M (1); 6.50 M; 7.40 M
47	2.38 S (3); 3.30 M (4); 4.20 M (1); 6.35 M (2); 7.30 M (11)

The compounds of formula I exhibit chemotherapeutic activity. In particular, they exhibit antimycotic activity, as indicated in vitro with tests against various families and types of mycetes, including *Trichophyton* 5 *quinckeanum*, *Aspergillus fumigatus*, *Microsporum canis*, *Sporotrichium schenkii* and *Candida albicans*, at concentrations of, for example 0.1 to 100 µg/ml, and in vivo in the experimental skin mycosis model in guinea pigs. In the latter model, guinea pigs are infected by cutaneous application of *Trichophyton quinckeanum*. The test substance is 10 administered daily for 7 days beginning 24 hours after the infection by local application by rubbing the test substance (taken up in polyethylene glycol) on the skin surface, or perorally, the test substance being administered as a 15 suspension. The activity is shown on local application at concentrations of from example 0.1 to 2%, in particular 0.1 to 0.6%. The oral activity is shown at dosages of, for example, 50 to 100 mg/kg.

The compounds are therefore indicated for use as 20 anti-mycotic agents. An indicated daily dose is from 500 to 2000 mg. If desired, this may be administered in divided doses 2 to 4 times a day in unit dosage form containing from about 125 mg to about 1000 mg or in sustained release form.

The compounds may be used in free base form or in the form of chemotherapeutically acceptable acid addition salts. Such salt forms exhibit the same order of activity as the free base forms.

5        The compounds may be admixed with conventional chemotherapeutically acceptable diluents and carriers, and, optionally, other excipients and administered in such forms as tablets or capsules. The compounds may alternatively be administered topically in such conventional forms as oint-  
10    ments or creams. The concentration of the active substance in such topical application forms will of course vary depending on the compound employed, the treatment desired and the nature of the form etc. In general, however, satisfactory results are obtained at concentrations of from 0.05 to 3.  
15    in particular 0.1 to 1 wt %.


A compound with particularly interesting activity is the compound of Example 4.

One group of compounds has a formula Ig,

0000896

900-9192





- 28 -



wherein  $R_1^I$  is 1-naphthyl, optionally mono-substituted  
by lower alkyl or alkoxy,

$u$  is a whole number from 1 to 8,

$R_6^{II}$  is of formula



5




wherein  $R_{19}$  is hydrogen, hydroxy, lower  
alkoxy or lower alkyl,

or of formula




wherein  $R_{20}$  is alkyl ( $\text{C}_{1-12}$ ) or phenylalkyl-  
( $\text{C}_{7-12}$ )

or of formula




wherein m, n and v are as defined above.

Another group of compounds comprises those of formula Ih,



wherein  $\text{R}_1^{\text{II}}$  is a radical of formula IIa, IIb wherein X is oxygen or sulphur, IIc, IId wherein s is 4, IIE wherein t is 3 or a radical of formula



wherein  $\text{R}_9$  is as defined above,

$\text{R}_2$  and  $\text{R}_5$  are independently hydrogen or lower alkyl,


u is a whole number from 1 to 8,

$\text{R}_6^{\text{III}}$  is as defined above for  $\text{R}_6$ ,


with the following provisos,

- (a)  $R_{10}$  is other than phenyl or phenylalkoxy, and
- (b) when  $R_1^{II}$  is 1-naphthyl optionally mono-substituted by lower alkyl or alkoxy and  $R_2$  and  $R_5$  are each hydrogen,  $R_6^{III}$  is other than
- 5 (i) a radical of formula IIIa, IIIb or IIIf,
- (ii) a radical of formula IIIc, IIId or IIIe,
- wherein  $R_{11}$ ,  $R_{12}$  and  $R_{13}$  are each hydrogen, or
- (iii) a radical of formula IIIg wherein one of  $R_{17}$  and  $R_{18}$  is hydrogen and the other is hydroxy, lower alkyl or lower alkoxy, or
- 10 (iv) an optionally substituted thiophen or furan radical.

A further group of formula I compounds comprises compounds of formula II,



- 15 wherein  $R_1^{III}$  is a radical of formula IIa, IIb wherein X is oxygen or sulphur, IIc, IIId wherein s is 4, IIe wherein t is 3, or a radical of formula






$R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_9$  are as defined above,  
with the proviso that  $R_3$  and  $R_4$  are other  
than  $-(CH_2)_4-$ ,

$R_6^{IV}$  is as defined above for  $R_6$  with respect to  
formula I, with the following provisos


5

- (i)  $R_{10}$  is other than phenyl or phenylalkoxy and
- (ii) when  $R_1$  is a radical of formula IIa,  $R_6^{IV}$  is  
other than a radical of formula IIIg, or  
phenyl.




and  $R_2$  represents hydrogen or lower alkyl,

or  $R_1$  and  $R_2$  together represent a group of formula



IIh



IIIi

whereby in the formulae IIa to IIIi,

$R_7$  and  $R_8$  represent, independently, hydrogen, halogen, tri-  
5 fluoromethyl, hydroxy, nitro, lower alkyl or lower alkoxy,

$R_9$  represents hydrogen, halogen, hydroxy, lower alkyl or  
lower alkoxy,

$X$  represents oxygen, sulphur, imino, lower alkyl imino or  
a radical of formula  $-(\text{CH}_2)_r^-$ ,

10  $p$  is 1, 2 or 3,

$r$  is 1, 2 or 3,

$s$  is 3, 4 or 5,

$t$  is 2, 3 or 4, and



$v$  is 3, 4, 5 or 6;

15  $R_3$  and  $R_5$  represent, independently, hydrogen or lower alkyl,  
and


$R_4$  represents  $C_{1-6}$  alkyl or  $C_{3-8}$  cycloalkyl- $(C_{1-6})$ -alkyl;

and

$R_6$  represents a group of formula



or



wherein  $R_{11}$  represents hydrogen, optionally  $\alpha$ -hydroxy substituted alkyl; alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, phenyl, phenalkyl or thienyl,

5

$R_{12}$ ,  $R_{13}$  and  $R_{14}$  represent, independently, hydrogen or lower alkyl, and

$\begin{array}{c} \text{=C} \\ \text{---} \end{array} \text{Z}$  represents a  $C_{5-8}$  cycloalkylidene radical optionally containing a double bond; or

10


b)  $R_1$  represents a group of formula IIa to IIg as defined under a),

$R_2$  represents hydrogen or lower alkyl,


$R_3$  and  $R_4$  together form a group  $-(CH_2)_u-$ , wherein  $u$  is an integer of 1 to 8, and

$R_5$  and  $R_6$  have the meanings given under a).


15 processes for their production, their use as pharmaceuticals and pharmaceutical compositions containing them.




IIb



IIc



IIId



IIe

wherein  $R_9$  is hydrogen, halogen of atomic number from 9 to 53, hydroxy, lower alkyl or lower alkoxy,

X is oxygen, sulphur, imino, lower alkylimino or a radical of formula


$-(CH_2)_r-$  wherein r is 1, 2 or 3,

s is 3, 4 or 5, and


t is 2, 3 or 4, and

$R_2$  is hydrogen or lower alkyl, or

(ii)  $R_1$  and  $R_2$  together with the carbon atom to which they are bound form a radical of formula IIIf or IIg,



IIIf




IIg

wherein p is 1, 2 or 3,

$R_3$  and  $R_5$ , independently, are hydrogen or lower alkyl,

$R_4$  is alkyl ( $C_{1-6}$ ), alkenyl ( $C_{3-12}$ ),  
alkynyl ( $C_{3-12}$ ) or cycloalkyl ( $C_{3-8}$ )-  
alkyl ( $C_{1-6}$ ); and

$R_6$  is (i) an aromatic, five-membered  
heterocycle containing one oxygen,  
sulphur or nitrogen hetero-ring atom  
and optionally an additional one or two  
nitrogen hetero-ring atoms and being  
optionally substituted on a carbon ring  
atom by halogen of atomic number from  
9 to 53, hydroxy, lower alkyl or lower  
alkoxy, and any nitrogen ring atom  
present being optionally substituted  
when possible, by lower alkyl, (ii) a  
radical of formula IIIa,




wherein  $R_9$  is as defined above,  
(iii) a radical of formula IIIb,



wherein  $R_{10}$  is alkyl ( $C_{1-12}$ ), alkenyl  
( $C_{3-12}$ ), alkynyl ( $C_{3-12}$ ) cycloalkyl ( $C_{3-8}$ )-  
alkyl ( $C_{1-6}$ ), phenyl-alkyl ( $C_{7-12}$ ),  
phenyl, phenylalkoxy ( $C_{7-16}$ ), or amino-  
alkyl ( $C_{1-12}$ );

(iv) a radical of formula IIIc, IIIId or IIIe,




wherein  $R_{11}$ ,  $R_{12}$  and  $R_{13}$ , independently, are hydrogen or lower alkyl,

$m$  is a whole number from 0 to 4,

$n$  is a whole number from 0 to 3, and

$v$  is a whole number from 0 to 5,

(v) a radical of formula IIIf,




wherein  $R_{14}$  is lower alkyl, alkoxy ( $C_{1-12}$ )-carbonyl, alkenyl ( $C_{3-12}$ ), alkynyl ( $C_{3-12}$ ), phenylalkyl ( $C_{7-12}$ ) or phenyl,

$R_{15}$  and  $R_{16}$ , independently, are hydrogen or lower alkyl, and

$q$  is a whole number from 0 to 5, or

(vi) a radical of formula IIIg



IIIg

wherein R<sub>17</sub> and R<sub>18</sub>, independently, are hydrogen, halogen of atomic number from 9 to 53, trifluoromethyl, hydroxy, nitro, lower alkyl or lower alkoxy, with the proviso that one of R<sub>17</sub> and R<sub>18</sub> is other than hydrogen, and with the general proviso that R<sub>1</sub> is not a radical of formula IIa when R<sub>6</sub> is a radical of formula IIIg or phenyl, R<sub>2</sub> is hydrogen and R<sub>3</sub> is hydrogen or lower alkyl,


b) R<sub>1</sub> is a radical of formula IIa to IIe, as defined above,

R<sub>2</sub>, R<sub>5</sub> and R<sub>6</sub> are as defined above, and R<sub>3</sub> and R<sub>4</sub> together are -(CH<sub>2</sub>)<sub>u</sub> - wherein u is a whole number from 1 to 8,


or an acid addition salt thereof.

2. A process for the production of a compound as claimed in Claim 1, which comprises

a) reacting a compound of formula IV,




wherein  $\text{R}_1$  to  $\text{R}_4$  are as defined above,  
with a compound of formula V,



wherein A is a leaving group, and

5  $\text{R}_5$  and  $\text{R}_6$  are as defined above, or


b) producing a compound of formula Ia,



wherein  $\text{R}_1$  to  $\text{R}_3$  and  $\text{R}_5$  are as defined above, and  
 $\text{R}_4^{\text{I}}$  and  $\text{R}_6^{\text{I}}$  are as defined above for  $\text{R}_4$  and  $\text{R}_6$   
respectively, with the proviso that they

10 each are other than alkynyl,

by reducing a compound of formula VI,






wherein  $R_1$  to  $R_3$ ,  $R_4^I$ ,  $R_5$  and  $R_6^I$  are as defined above,


or

c) producing a compound of formula Ib,




wherein  $R_1$  to  $R_6$  are as defined above,

5 by isomerising photochemically a compound of formula Ic,



wherein  $R_1$  to  $R_6$  are as defined above, or

d) producing a compound of formula Id,




wherein  $R_1$ ,  $R_2$ ,  $R_5$  and  $R_6$  are as defined above,

$R_3^I$  is hydrogen or lower alkyl,

$R_4^{II}$  is alkyl( $C_{1-6}$ ), alkenyl( $C_{3-12}$ ), alkynyl


( $C_{3-12}$ ) or cycloalkyl( $C_{3-8}$ )alkyl( $C_{1-6}$ );

10 by introducing the group  $R_4^{II}$  into a compound of formula VII,



wherein  $R_1$ ,  $R_2$ ,  $R_3^I$ ,  $R_5$  and  $R_6$  are as defined above.





3. A compound of Claim 1, having the formula Ig,



wherein  $R_1^I$  is 1-naphthyl, optionally mono-substituted  
by lower alkyl or alkoxy,

5  $u$  is a whole number from 1 to 8,



$R_6^{II}$  is of formula



$$-\text{CO}-\text{OR}_{20} \quad (\text{e})$$


wherein  $R_{20}$  is alkyl ( $C_{1-12}$ ) or phenylalkyl-  
( $C_{7-12}$ )

or of formula




wherein m, n and v are as defined in  
Claim 1.

4. A compound of Claim 1, having the formula Ia,



wherein  $R_1^{II}$  is a radical of formula IIa, IIb wherein X is oxygen or sulphur, IIc, IIId wherein s is 4, IIe wherein t is 3 or a radical of formula



wherein  $R_9$  is as defined in Claim 1,  
 $R_2$  and  $R_5$  are independently hydrogen or lower  
 alkyl,


$u$  is a whole number from 1 to 8,

$R_6^{III}$  is as defined in Claim 1 for  $R_6$ ,


with the following provisos,

- (a)  $R_{10}$  is other than phenyl or phenylalkoxy, and
- (b) when  $R_1^{II}$  is 1-naphthyl optionally mono-substituted by  
 lower alkyl or alkoxy and  $R_2$  and  $R_5$  are each hydrogen,  
 $R_6^{III}$  is other than
  - (i) a radical of formula IIIa, IIIb or IIIf,
  - (ii) a radical of formula IIIc, IIId or IIIe,  
 wherein  $R_{11}$ ,  $R_{12}$  and  $R_{13}$  are each hydrogen, or
  - (iii) a radical of formula IIIg wherein one of  $R_{17}$   
 and  $R_{18}$  is hydrogen and the other is hydroxy,  
 lower alkyl or lower alkoxy, or
  - (iv) an optionally substituted thiophen or furan  
 radical.

5. A compound of Claim 1 having the formula Ii,



wherein  $R_1^{III}$  is a radical of formula IIa, IIb wherein X is oxygen or sulphur, IIc, IId wherein s is 4, IIe wherein t is 3, or a radical of formula



5

$R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_9$  are as defined in Claim 1, with the proviso that  $R_3$  and  $R_4$  are other than  $-(CH_2)_u-$ ,

5

$R_6^{IV}$  is as defined in Claim 1 for  $R_6$  with respect to formula I, with the following provisos

10

- (i)  $R_{10}$  is other than phenyl or phenylalkoxy and
- (ii) when  $R_1$  is a radical of formula IIa,  $R_6^{IV}$  is other than a radical of formula IIIg, or phenyl:

0000896

900-9192

- 12 -

6. A compound of Claim 1 which is N-(3-cyclohexyl-2-trans-propenyl)-N-methyl-N-(1-naphthylmethyl)amine.

7. A pharmaceutical composition comprising a compound of any one of Claims 1, and 3 to 6 in free base  
5 form or in chemotherapeutically acceptable acid addition salt form in association with a chemotherapeutically acceptable diluent or carrier.

3700/RR/HD